

The Kondo screening cloud: what can we learn from perturbation theory?

Victor Barzykin¹ and Ian Affleck^{1,2}

*Department of Physics¹ and Canadian Institute for Advanced Research²,
University of British Columbia, Vancouver, BC, V6T 1Z1, Canada*

We analyse the role which the distance scale $\xi_K = v_F/T_K$ plays in the single-impurity Kondo problem using renormalization group improved perturbation theory. We derive the scaling functions for the local spin susceptibility in various limiting cases. In particular, we demonstrate exactly that the non-oscillating part of it should be short-range, i.e., vanish for distances $r \gg 1/k_F$ and show explicitly that the interior of the screening cloud *does not* exhibit weak coupling behavior.

PACS numbers: 75.20.Hr, 75.30.Mb, 75.40.Cx

Although the Kondo effect has been very thoroughly studied for over thirty years [1], comparatively little theoretical work concentrated on the role of spatial correlations. Part of the problem is that it is inaccessible to the Bethe ansatz and difficult to analyse using Wilson's renormalization group. No theoretical and experimental consensus has emerged on the question of the length scale at which screening of the impurity takes place. On one hand, in the scaling language [2,3], the low energy scale $T_K \propto e^{-1/\rho J}$ implies the presence of the exponentially large length scale $\xi_K = v_F/T_K$; here v_F is the Fermi velocity, T_K is the Kondo temperature, J is the Kondo coupling, and ρ is the density of states (per spin). On the other hand, in the Knight shift experiments of Boyce and Slichter [4] no such scale was observed. It also appears experimentally [1] that alloys with impurity concentration $n \gg (1/\xi_K)^3$, i.e., where the inter-impurity distances are much less than ξ_K , display single-impurity behavior. In this letter we attempt to clarify these questions using renormalization group (RG) improved perturbation theory. Related theoretical work includes early perturbative calculations [5], and RG approaches [6,7]. Recent theoretical work [8,9] has also addressed these issues.

In what follows we consider the standard $S_{\text{imp}} = 1/2$ Kondo model,

$$H = v_F \int_{-\infty}^{\infty} dr \psi_L^\dagger(r) (id/dr) \psi_L(r) + v_F \lambda \psi_L^\dagger(0) \frac{\sigma}{2} \psi_L(0) \cdot \mathbf{S}_{\text{imp}}. \quad (4)$$

The local spin susceptibility $\chi(r, T)$ in this formalism is a sum of uniform and $2k_F$ parts.

$$\chi_{un}(r, T) \equiv (v_F/T) \langle [\psi_L^\dagger(r) \frac{\sigma^z}{2} \psi_L(r) + \psi_L^\dagger(-r) \frac{\sigma^z}{2} \psi_L(-r)] S_{\text{tot}}^z \rangle, \quad (5)$$

and χ_{2k_F} is given by the same expression with r replaced by $-r$ in the argument of ψ_L . Here $\mathbf{S}_{\text{tot}}$ is the total spin in the one-dimensional theory:

$$\begin{aligned} \mathbf{S}_{\text{tot}} &\equiv \mathbf{S}_{\text{imp}} + \mathbf{S}_{\text{el}} \\ \mathbf{S}_{\text{el}} &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dr \psi_L^\dagger(r) \frac{\sigma}{2} \psi_L(r). \end{aligned} \quad (6)$$

$$H = \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} \psi_{\mathbf{k}}^{\dagger\alpha} \psi_{\mathbf{k}\alpha} + J \mathbf{S}_{\text{imp}} \cdot \sum_{\mathbf{k}, \mathbf{k}'} \psi_{\mathbf{k}}^{\dagger\alpha} \frac{\sigma_{\alpha}^{\beta}}{2} \psi_{\mathbf{k}'\beta}. \quad (1)$$

The quantity measured in the Knight shift experiments is the local spin susceptibility,

$$\chi(r, T) \equiv (1/T) \langle \psi^\dagger(\mathbf{r}) \frac{\sigma^z}{2} \psi(\mathbf{r}) S_{\text{tot}}^z \rangle - \chi_0, \quad (2)$$

where $S_{\text{tot}}^z = S_{\text{imp}}^z + (1/2) \int d\mathbf{r} \psi^\dagger(\mathbf{r}) \sigma^z \psi(\mathbf{r})$ is the total spin operator of the impurity and conduction electrons. The bulk Pauli contribution, $\chi_0 \approx \rho/2$ has been subtracted. Recently a scaling conjecture was made [8], supported by numerical results, that in the scaling limit, $rk_F \gg 1$, $T \ll E_F$, the spin susceptibility has the following form:

$$\chi = \frac{\chi_{2k_F} \left(\frac{rT}{v_F}, \frac{T}{T_K} \right)}{4\pi^2 r^2 v_F} \cos(2k_F r) + \frac{\chi_{un} \left(\frac{rT}{v_F}, \frac{T}{T_K} \right)}{8\pi^2 r^2 v_F}, \quad (3)$$

where χ_{2k_F} and χ_{un} are universal functions of two scaling variables [10]. This form follows from the relativistic one-dimensional formulation of the Kondo problem [11]. The one-dimensional Hamiltonian in terms of the left-moving fields is:

Renormalizability implies that the functions χ_A ($A = 2k_F, un$) obey equations of the form:

$$\left[D \frac{\partial}{\partial D} + \beta(\lambda) \frac{\partial}{\partial \lambda} + \gamma_A(\lambda) \right] \chi_A(T, \lambda, D, rT/v_F) = 0, \quad (7)$$

where D is the ultra-violet cut-off (the bandwidth), $\lambda \equiv \rho J$ is the dimensionless coupling constant, $\beta(\lambda)$ is the β -

function and $\gamma_A(\lambda)$ is the anomalous dimension, which is a sum of contributions from the local fermion bilinears and from S_{tot}^z . In this case, both contributions to γ_A vanish. S_{tot}^z has zero anomalous dimension because it is a conserved operator. The fermion bilinear at $r \neq 0$ has vanishing anomalous dimension because the interactions occur only at the origin; only “boundary operators” receive anomalous dimensions in such theories [11]. The most general solution of this equation is a general function of the effective coupling constant at scale T , λ_T and of the parameter rT/v_F . Swapping the dependence on λ_T for dependence on T/T_K and using the fact that χ_{2k_F} is real [8] we obtain Eq. (3).




FIG. 1. Cancellation of the uniform part of the local spin susceptibility.

Consider first χ_{un} . From our lower-order perturbative analysis we have found that χ_{un} is zero. This fact is indeed quite general, and can be proven to all orders in perturbation theory. We will denote as $\chi_{un, \text{imp}}(r)$ the part of the uniform spin susceptibility for which the im-

purity piece S_{imp} of S_{tot} is responsible, and the other part $\chi_{un, \text{el}}(r)$. [These two contributions are defined as zero frequency Fourier transforms. Their sum involves a conserved operator, S_{imp}^z , and so Fourier transforming was equivalent to dividing by T in Eq. (2).] The fact that $\chi_{un, \text{imp}}(r)$ vanishes can very easily be seen. In all orders of perturbation theory, one integrates over two free electron Green's functions, G , connecting the points (r, τ) and $(0, \tau_i)$ (see Fig. 1a). The result of the τ -integration is zero:

$$I = \int_0^\beta d\tau G(r, \tau - \tau_1) G(-r, \tau_2 - \tau) \quad (8)$$

$$= \int_0^\beta \frac{d\tau (\pi T)^2}{\sin[\pi T(\tau - \tau_1 + ir)] \sin[\pi T(\tau_2 - \tau - ir)]} = 0,$$

because after the change of integration variable, $\tau \rightarrow \exp(i2\pi T\tau)$, one encounters contour integration with two poles on one side.

The cancellation of $\chi_{un, \text{el}}(r)$ is somewhat less trivial, since in addition to the graphs with this type of integration, there are other graphs. It can be shown that these graphs sum up to a graph where the integration of the above type is present (see Fig. 1b) and therefore also vanish. As a result, $\chi_{un}(r, T)$ exactly vanishes at finite r . This function, however, may include a singular contribution at $r = 0$ (or, more correctly, at r of the order the ultraviolet cut-off in the one-dimensional theory, which is essentially $1/k_F$).

We will now analyse the $2k_F$ part of the spin susceptibility $\chi_{2k_F}(r, T)$. From our analysis of the perturbation theory up to third order, we find that the scaling form Eq.(3) is indeed obeyed. The scaling function χ_{2k_F} can be written:

$$\chi_{2k_F} \left(x = \frac{rT}{v_F}, \lambda_T \right) = \frac{(\lambda_E + (3\pi/2)\lambda_E^2 x + \text{const}\lambda_E^3)(1 - \lambda_T)}{(4/\pi^2) \sinh(2\pi x)}, \quad (9)$$

where λ_T is given by:

$$\lambda_T = \lambda + \lambda^2 \ln(D/T) + \lambda^3 [\ln^2(D/T) - (1/2) \ln(D/T) + \text{constant}]. \quad (10)$$

λ_E is also given by Eq.(10), with T replaced by another effective scale, $E(x) = T/[1 - \exp(-4\pi x)]e$. Eq. (10) is universal up to a rescaling of the cut-off, D and a change in the *constant* term. We use this freedom to redefine D to simplify our expressions. It is important to note that the infrared divergences of perturbation theory are *not* cut off at low T by going to small r , as was first noticed by Gan [7]. It is also necessary to have a high T so that λ_T is small. In the third order, these divergences are associated with the graph shown in Fig. 2. Due to the non-conservation of momentum by the Kondo interaction, the bubble on the right gives a logarithmic T -dependent factor which is independent of r . Thus, at

low T , the interior of the screening cloud does *not* exhibit weak coupling behavior. $\chi(r, T)$ decreases exponentially, $\propto e^{-2\pi x}$, for $r \gg v_F/T$. In the opposite limit, $r \ll v_F/T$, χ_{2k_F} can be expressed as a polynomial in λ_T and $\ln x$, or equivalently λ_T and λ_r , the effective coupling at scale r . (Note that $\lambda_r \equiv \lambda_E$ for $r \ll v_F/T$.) To third order, the local spin susceptibility has the following form:

$$\chi_{2k_F}(\lambda_T, \lambda_r) = \frac{(\lambda_r + c\lambda_r^3)}{(r/\pi v_F)} \frac{(1 - \lambda_T)}{4T} \quad (x \ll 1), \quad (11)$$

where c is a constant. The T -dependent factor is, to the order we work, precisely the total impurity susceptibility, $\chi_{\text{tt}}(T)$. This is the total susceptibility less the bulk

Pauli term and its value has been determined accurately [1]. At low T it approaches $1/T_K$. $\lambda_r \ll 1$ provided that $r \ll \xi_K$, even if $T \ll T_K$.




FIG. 2. Singular third-order graph for $\chi(r, T)$.

Boyce and Slichter [4] have measured the Knight shift from Cu nuclei near the doped Fe impurities, at distances up to 5-th nearest neighbor. At these very small distances of order of a few lattice spacings, they have found empirically that the Knight shift obeyed a factorized form, $\chi(r, T) \approx f(r)/(T + T_K)$, with rapidly oscillating function $f(r)$ for a wide range of T extending from well above to well below the Kondo temperature. This is essentially the same form as Eq. (11) at $T \gg T_K$ and low r . We observe from Eq.(9) that the factorization breaks down for $r > v_F/T$. Our perturbative approach isn't valid unless $T \gg T_K$, so we can't check factorization at low T . This question can certainly be addressed for the overscreened large- k Kondo problem, where it is possible to obtain reliable low-temperature results from the weak-coupling perturbative expansion [12]. The factorized behavior of the local spin susceptibility was also obtained in Ref. [13].

$$4T\chi_j(T, \lambda, \Lambda) = \exp \left[\int_{\lambda}^{\Lambda} \frac{\gamma_j(\lambda')}{\beta(\lambda')} d\lambda' \right] \Pi_j(\lambda_T) = \Phi_j(\lambda_T) \exp \left[- \int_0^{\lambda} \frac{\gamma_j(\lambda')}{\beta(\lambda')} d\lambda' \right]. \quad (14)$$

Here $\Phi_j(\lambda_T)$, $\Pi_j(\lambda_T)$ are some scaling functions. From our third-order perturbative analysis using Wilson's result [2] for $\chi_{tt}(T)$ we have obtained that the functions $\Phi_j(\lambda_T)$ coincide for all three susceptibilities up to and including terms of order λ_T^2 . When λ is small, the scale factor in Eq.(14) can be easily calculated from perturbative expansion of β and γ_{imp} :

$$\exp \left[- \int_0^{\lambda} \frac{\gamma_{\text{imp}}(\lambda')}{\beta(\lambda')} d\lambda' \right] \simeq 1 + \frac{\lambda}{2}. \quad (15)$$

Thus, at least at high temperatures, where our perturbative calculation of the scaling functions is valid, the integrated electronic susceptibility obeys:

$$\int \chi(r, T) dr \approx -\frac{\lambda}{2} \chi_{tt}(T). \quad (16)$$

This result is mostly given by the electron-impurity correlator, while the electron-electron piece,

$$\chi_{ee} \equiv \int_0^{\beta} d\tau \langle S_{el}^z(\tau) S_{el}^z(0) \rangle \simeq \frac{\lambda^2}{4} \chi_{tt}(T), \quad (17)$$

So far, we have only discussed the region $r \gg 1/k_F$. It is also interesting to consider the integral of $\chi(\vec{r})$ over all space. Although the $2k_F$ oscillations tend to cancel at large r , there could be a non-zero contribution from $r \leq 1/k_F$. It is useful to consider various static spin susceptibilities:

$$\chi_{tt}(T) \equiv \frac{\langle S_{\text{tot}}^z S_{\text{tot}}^z \rangle}{T}, \quad \chi_{ti}(T) \equiv \frac{\langle S_{\text{imp}}^z S_{\text{tot}}^z \rangle}{T},$$

$$\chi_{ii}(T) \equiv \int_0^{\beta} \langle S_{\text{imp}}^z(\tau) S_{\text{imp}}^z(0) \rangle d\tau. \quad (12)$$

It is easy to see that the electron spin polarization in the presence of an impurity is determined by the spatial integral of Eq.(2), or, equivalently, by $\chi_{tt}(T) - \chi_{ti}(T)$. Since S_{tot}^z is conserved, these susceptibilities obey the RG equation, Eq. (7), with anomalous dimensions determined by the anomalous dimension $\gamma_{\text{imp}}(\lambda)$ of the operator S_{imp}^z . For the three different susceptibilities: $\gamma_{tt} = 0$, $\gamma_{ti} = \gamma_{\text{imp}}$, and $\gamma_{ii} = 2\gamma_{\text{imp}}$. The low-order perturbative results for $\beta(\lambda)$ [14] and $\gamma_{\text{imp}}(\lambda)$ [7] are:

$$\beta(\lambda) = -\lambda^2 + \frac{\lambda^3}{2}, \quad \gamma_{\text{imp}}(\lambda) = \frac{\lambda^2}{2}. \quad (13)$$

We should note that the RG equations are only simple for the choice of spin correlators Eq.(12). The equations become mixed if written in terms of impurity, impurity-electron, and electron parts of the spin susceptibility. The general solution of these RG equations has the form:

is further suppressed by a power of λ . In the scaling limit of small bare coupling, $\lambda \rightarrow 0$, the total polarization of the conduction electrons vanishes, at least at high temperature. The oscillating function $\chi(r)$ integrates to 0, and χ_{tt} comes entirely from the impurity-impurity part. It should be emphasized, however, that the result is non-zero at finite bare coupling λ . (A typical experimental value of λ might be $1/\ln(E_F/T_K) \approx .15$.) We conjecture that the equality of the scaling functions, $\Phi_j(\lambda_T)$ defined in Eq. (14) holds at all T , so that Eq. (16) is true at all T and small bare coupling. In particular, the integrated electronic susceptibility then vanishes in the scaling limit of zero bare coupling at all T . Precisely this result was found at $T = 0$ from the Bethe ansatz [15]. However, this conjecture is *not* completely consistent with recent work of Lesage et al. [16] which extrapolates to the isotropic Kondo Hamiltonian from an anisotropic model. While this may well indicate that our conjecture is wrong, it is

also possible that there is a problem with the extrapolation since we find the susceptibilities to be very singular in the isotropic limit.

$\chi_{2k_F}(r, T)$ is a sum of impurity and electron parts, $\chi_{2k_F, \text{imp}}$ and $\chi_{2k_F, \text{el}}$. While the former obeys the RG equation Eq.(7) with $\gamma_A = \gamma_{\text{imp}}$, the latter obeys a more complicated RG equation due to operator mixing of $\mathbf{S}_{\text{el}}$ and $\mathbf{S}_{\text{imp}}$. $\chi_{2k_F, \text{imp}}$ has the same λ -dependent factor as in Eq. (15) multiplied by a scaling function:

$$\chi_{2k_F, \text{imp}} \simeq \left(1 + \frac{\lambda}{2}\right) \chi_{2k_F}^{(1)}(\lambda_T, x), \quad (18)$$

where the scaling function

$$\chi_{2k_F}^{(1)}(\lambda_T, x) = \frac{(\lambda_E + \text{const} \lambda_E^3)(1 - \lambda_T)}{(4/\pi^2) \sinh(2\pi x)} \quad (19)$$

is not the same as in Eq.(9). Thus, we obtain for all r at weak bare coupling:

$$\chi_{2k_F, \text{el}} \approx -\frac{\lambda}{2} \chi_{2k_F}^{(1)}(\lambda_T, x) + \frac{(3\pi/2)\lambda_E^2 x(1 - \lambda_T)}{(4/\pi^2) \sinh(2\pi x)}. \quad (20)$$

Hence two different scaling functions are present in the experimentally measured Knight shift, and their share depends upon the gyromagnetic ratios for the impurity and conduction electrons. Unlike for the total spin susceptibility, $\chi_{2k_F, \text{el}}$ doesn't vanish in the scaling limit of zero bare coupling. However, it is small compared to $\chi_{2k_F, \text{imp}}$ when the bare coupling and x are both small.

We may summarize the response of the weak-coupling Kondo model to a small magnetic field as follows. The impurity spin is much more strongly affected by a weak Kondo coupling than is the electron gas. This is connected with the fact that the free impurity susceptibility blows up as $T \rightarrow 0$, whereas the free conduction electron susceptibility does not. Thus even a weak Kondo coupling drastically affects χ_{tt} at low T , causing the diverging Curie susceptibility to level off at $1/T_K$. On the other hand the affect on the electron gas can only become appreciable in a long distance, infrared limit. The effect of this, however, is not very dramatic, because of the factor of $1/r^2$, which arises for purely dimensional reasons.

At short distances, of $O(1/k_F)$, the excess polarization of the electron gas produced by the Kondo interaction is small. This together with the oscillating nature of the long-distance polarization gives rise to a small integrated excess polarization of the electron gas of $O(\lambda/T_K)$.

We would like to thank J. Gan, F. Lesage, N. Prokof'ev, H. Saleur, E. S. Sørensen, P. C. E. Stamp, C. Varma and A. Zawadowski for useful discussions and comments. This research was supported by NSERC of Canada.

-
- [1] A. C. Hewson, *The Kondo Problem to Heavy Fermions*, Cambridge University Press, Cambridge 1993.
 - [2] K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975).
 - [3] (a) Ph. Nozières, J. Low Temp. Phys. **17**, 31 (1974); Ph. Nozières, J. de Phys. **39**, 1117 (1978); (b) For a review see Ph. Nozières, in *Proc. 14th Int. Conf. on Low Temperature Physics*, eds. M. Krusius and M. Vuorio, Vol. 5, (North Holland, Amsterdam 1975).
 - [4] J. P. Boyce and C. P. Slichter, Phys. Rev. Lett. **32**, 61 (1974); Phys. Rev. B **13**, 379 (1976)
 - [5] J. Kondo, Solid State Physics **23**, 183 (1969).
 - [6] K. Chen, C. Jayaprakash and, H. R. Krishnamurthy, Phys. Rev. B **45**, 5368 (1992).
 - [7] J. Gan, J. Phys.:Cond. Mat. **6**, 4547 (1994).
 - [8] E. S. Sørensen and I. Affleck, preprint (1995), to appear in Phys. Rev. B; cond-mat 9508030.
 - [9] O. Újsághy, A. Zawadowski, and B. Gyorffy, preprint 1995.
 - [10] Definitions of χ_{2k_F} and χ_{un} differ by a factor of v_F from those used in Ref. [8].
 - [11] See, for example, I. Affleck and A. W. W. Ludwig, Nucl. Phys. B **360**, 641 (1991).
 - [12] V. Barzykin, I. Affleck, in progress
 - [13] E. Kim, M.S. Makivik, D.L. Cox, Phys. Rev. Lett. **75**, 2015 (1995).
 - [14] A. A. Abrikosov and A. A. Migdal, J. Low. Temp. Phys. **3**, 519 (1970); M. Fowler and A. Zawadowski, Sol. State Comm. **9**, 471 (1971).
 - [15] J. H. Lowenstein, Phys. Rev. B **29**, 4120.
 - [16] F. Lesage, H. Saleur and S. Skorik, cond-mat/9603043.